Dietary *n*-6 and *n*-3 fatty acids in immunity and autoimmune disease

Laurence S. Harbige

School of Chemical and Life Sciences, University of Greenwich, Wellington Street, London SE18 6PF, UK

Lipid biology plays a significant role in the normal and pathological functioning of cells of the immune system (Hopkins et al. 1981; Dvorak et al. 1983; Schlager et al. 1983; Shipman et al. 1988; Roper et al. 1990; Goetzl et al. 1995; Pushkareva et al. 1995; Hennig et al. 1996). In addition to the physiological requirements of cells of the immune system for essential fatty acids, dietary fatty acid modulation of the membrane composition and functions of immune cells can affect both normal and pathological processes (Calder, 1996; Harbige, 1996; Hennig et al. 1996). Among the first immunologists to recognize the importance of dietary fatty acids in immunity and autoimmune disease were Peter Medawar and Jürgen Mertin (Medawar et al. 1979) in the UK and Gabriel Fernandes and Robert Good (Fernandes et al. 1972) in the USA. The importance of dietary fatty acids in the prevention or control of autoimmune disease in animal models is now well documented (Mertin, 1981; Levy et al. 1982; Kelley et al. 1985; Morrow et al. 1985; Robinson et al. 1986; Alexander et al. 1987; Harbige et al. 1995; Lin et al. 1996). However, the biochemical and immunological mechanisms by which dietary fatty acids affect the immune system and autoimmunity have remained largely obscure. In recent years important new concepts in immunoregulation have emerged which are likely to be of significance to effects of fatty acid nutrition on immunity. Mosman & Coffman (1989) have characterized two different patterns of cytokine secretion by T-cells which lead to different functional responses. T helper₁ (Th₁) cells produce interleukin (IL)-2, interferon- γ and tumour necrosis factor- β which are not synthesized by Th₂ cells. In contrast, Th₂ cells (but not Th₁) produce IL-4, IL-5 and IL-10. Th₁ cells enhance cell-mediated inflammatory activity, whereas Th₂ cells synthesize cytokines that help B-cells develop into antibody-producing cells. There are also T-cells able to produce both Th₁ and Th₂ cytokines, referred to as Th₀ cells (Male et al. 1996). Weiner (1997) has further characterized a Th₃ T-cell subset that primarily produces transforming growth factor- β (TGFB), provides help for immunoglobulin (Ig) A production, and has suppressive properties. Importantly, the overall balance of cytokine production by Th₁, Th₂, and Th₃ cells affects the type of immune response generated, e.g. Th₂ cytokines can down-regulate production of Th₁ cytokines and vice versa (Male et al. 1996). Eicosanoids are known to affect these immunoregulatory mechanisms (Phipps et al. 1991), which may, in part, explain the complex relationship between the immune system, eicosanoids and the eicosanoid precursor polyunsaturated fatty acids derived from the diet. The balance of membrane fatty acids and eicosanoids is also known to influence inflammatory reactions (Terano *et al.* 1984; Lefkowith, 1988; Tate *et al.* 1989; Ross, 1993; Calder, 1996; Hennig *et al.* 1996).

Essential fatty acid-deficient diets and high- and low-fat diets in autoimmune disease

Diets deficient in essential fatty acids and diets low in fat markedly increase the survival and reduce spontaneous autoimmune disease in NZB \times NZW F₁ mice (primarily autoantibody mediated), a model of the human disease systemic lupus erythematosus (Hurd et al. 1981; Levy et al. 1982). Essential fatty acid-deficient diets also protect against autoimmune diabetes in the BB rat and in a low-dosestreptozotocin-treated mouse model of autoimmune diabetes (Lefkowith et al. 1990). High-fat (both lard and maize oil) diets increase the levels of natural thymocytotoxic and anti-DNA antibodies along with increased immune complex deposition, decrease T-cell mitogenic responses to concanavalin A (Con A), and accelerate the disease course in NZB × NZW F_1 mice (Levy et al. 1982; Yumura et al. 1985). In addition, the same group (Morrow et al. 1985) reported that high-saturated-fat (lard) diets have deleterious effects on both macrophage phagocytosis and natural killer cell activity, the latter correlating with in vitro interferon production. In the NZB \times NZW F₁ autoimmune model a high-fat diet consisting of equal amounts of lard and soyabean oil (rich in linoleic acid) causes animals to develop more severe disease and have a shortened lifespan associated with increased IgG anti-cardiolipin antibodies (Lin et al. 1996, 1997). In contrast to some of the previously described findings, in rats clinical manifestations of a T-cell-mediated disease experimental autoimmune encephalomyelitis (EAE), an animal model for the human disease multiple sclerosis, appear potentiated by fat deficiency (Clausen & Moller, 1967; Selivonchick & Johnston, 1975). However, there does appear to be disparity in the effects of fat deficiency on EAE (Levine & Sowinski, 1980), and the effects of high-fat diets have not been investigated.

Abbreviations: Con A, concanavalin A; EAE, experimental autoimmune encephalomyelitis; Ig, immunoglobulin; IL, interleukin; LT, leukotriene; ova, ovalbumin; PG, prostaglandin; PHA, phytohaemagglutinin; Th, T helper; TGFβ, transforming growth factor-β.
Corresponding author: Dr L. S. Harbige, fax +44 (0)181 331 8305, email L. Harbige@greenwich.ac.uk

Autoimmunity, immune functions and *n*-3 fatty acids

Animal and human studies

The effects of dietary n-3 fatty acids on ex vivo lymphocyte functions, as judged by mitogen stimulation, have been consistent in both human and animal studies showing suppressed responses (Calder, 1996). Kelley et al. (1988) found that an n-3 fatty acid-rich diet (76 g fish oil/kg) decreased rabbit peripheral-blood lymphocyte mitogenic responses to Con A, phytohaemagglutinin (PHA) and pokeweed (Phytolacca americana) mitogen ex vivo. Similarly, in rats fed on a high-fish-oil diet Yaqoob et al. (1994) reported decreased proliferative responses to Con A for spleen and lymph node lymphocytes. In mice fed on a high-fish-oil diet spleen lymphocytes showed decreased responses to Con A (Yaqoob & Calder, 1995). In both fish oil-fed and fish oil plus highor low-dose oral ovalbumin (ova) tolerance induction, ova immunized mice exhibited markedly reduced spleen lymphocyte responses to ova ex vivo and, in ova-fed animals only, reduced ova-specific levels of serum IgG₁, IgG_{2a} and IgG_{2b} (LS Harbige and BAC Fisher, unpublished results). In contrast, Morrissey et al. (1990) observed increased Con A -induced proliferation with a low-fish-oil diet (50 g/kg) plus additional vitamin E (150 mg/kg). This suggests that low-dose-fish-oil feeding may have different effects on proliferative responses, or that vitamin E may have an effect on mitogen-induced lymphoproliferation independent of fish oil n-3 fatty acids. Interestingly, Kelley et al. (1988) observed enhanced lymphocyte responses in rabbits fed on α -linolenic acid-rich linseed oil. However, this response may reflect evolutionary adaptation of rabbits to an α -linolenic acid-rich herbivorous diet, and highlights the importance of species differences. All the previously described ex vivo findings may not reflect fully those in vivo. For example, there appear to be site-specific differences in lymphocyte responses to dietary fatty acids (Yaqoob et al. 1994), which appear to be dependent on the local adipose-tissue lipid composition (Pond, 1996). It is interesting to note that adipose tissue around lymph nodes preferentially incorporates or selectively retains polyunsaturated fatty acids (Pond, 1996). This suggests that lymphoid cells inside lymph nodes can be locally supplied with the appropriate fatty acids for membrane and eicosanoid synthesis.

Several investigators have found reduced major histocompatibility complex class II molecule expression in rodents fed on fish oil (Huang et al. 1992; Sherrington et al. 1995), indicating a decreased ability of antigen-presenting cells to present antigen to T lymphocytes. Consistent with the previous findings, Fujikawa et al. (1992) found splenocytes acting as antigen-presenting cells from mice fed on the n-3fatty acid eicosapentaenoic acid had a reduced ability to present antigen to T-cell clones. Experimental data on the effects of feeding n-3 fatty acids on macrophage phagocytic functions and the production of H₂O₂, superoxide and NO have been inconsistent (Calder, 1996). There are also inconsistent reports of the effects of feeding fish oil to rodents on tumour necrosis factor and IL-1 production by peritoneal macrophages (Calder, 1996). In rats fed on fish oil (Leitch et al. 1984) or pure ethyl ester of eicosapentaenoic acid (Terano et al. 1984), ex vivo peritoneal leucocytes produced

less pro-inflammatory leukotriene (LT) B_4 and more of the less-potent LTB₅.

Hughes et al. (1995) found that fish oil supplementation in human subjects reduced expression of major histocompatibility complex class II DR molecules on peripheral-blood monocytes which suggests antigen presentation may be affected. Fish oil supplementation for 24 weeks reduced the relative percentage of peripheral blood CD4+ and increased the percentage of CD8+ cells in man (Meydani et al. 1993). In non-human primates fed on diets containing 1.3 and 3.3 % energy as eicosapentaenoic acid and docosahexaenoic acid for 14 weeks increased peripheral-blood lymphocyte responses to Con A and PHA have been reported (Wu et al. 1996). These investigators also found that the plasma vitamin E level was maintained, and suggested that in the presence of adequate vitamin E concentrations lymphocyte mitogenic proliferative responses are enhanced. Again this indicates vitamin E may have independent effects on immune function and that the balance of n-3 fatty acids and vitamin E is important. The effects of fish oil supplementation have been extensively studied on human peripheral-blood mononuclear cell cytokine production ex vivo, particularly IL-1 α , IL-1 β , IL-2, IL-6, and tumour necrosis factor- α , showing decreased production of these cytokines (Calder, 1996). Administration of 12-Otetradecanoylphorbol-13-acetate induces tumour necrosis factor- α gene expression in human HL-60 cells through arachidonic acid metabolites acting as regulating messengers (Horiguchi et al. 1989). It is well known that on feeding fish oil the n-3 fatty acid, eicosapentaenoic acid, displaces arachidonic acid in membrane phospholipids. It is possible, therefore, that the effects of n-3 fatty acids on some proinflammatory cytokines may be indirect, and may indicate the direct importance of arachidonic acid metabolism in their regulation. The delayed-type hypersensitivity response, which is an in vivo measure of cell-mediated immunity, was reported by Meydani et al. (1993) to be reduced in response to several recall antigens, including tetanus toxoid and Mycobacterium tuberculosis, in subjects supplemented daily with 1.23 g eicosapentaenoic acid and docosahexaenoic acid. Investigators have shown that feeding fish oil induces a decrease in pro-inflammatory neutrophil LTB₄ production and an increase in the less potent pro-inflammatory lipidmediator LTB₅ in human subjects with inflammatory diseases (Schmidt & Dyerberg, 1989).

Autoimmune disease

In mouse models of the human disease, systemic lupus erythematosus, fish oils rich in *n*-3 fatty acids increase the longevity and delay the onset of clinical manifestations of spontaneous autoimmunity (autoantibody mediated) in NZB × NZW F₁ and MRL/Ipr mice (Kelley *et al.* 1985; Godfrey *et al.* 1986; Robinson *et al.* 1986). Feeding fish oil compared with maize oil delayed the onset of autoimmune disease in NZB × NZW F₁ mice (Fernandes *et al.* 1994). This observation was associated with elevated IL-2, IL-4 and TGF β_1 and lower c-myc and c-ras mRNA in the spleen, with protein products following the same pattern (Fernandes *et al.* 1994). The same group reported high kidney TGF β_1 mRNA for NZB \times NZW F₁ mice fed on maize oil and low kidney TGF β_1 mRNA expression for NZB \times NZW F₁ mice fed on fish oil, the reverse of their findings in the spleen (Fernandes, 1994). In contrast, antioxidant enzyme mRNA expression in the liver was higher in fish oil-fed mice than in maize oil-fed mice (Fernandes, 1994). These workers suggested that these findings were important to the renal disease normally observed in the NZB \times NZW F₁ mouse, and also that dietary fatty acids regulate $TGF\beta_1$ and antioxidant enzyme expression in an organ-specific manner (Chandrasekar et al. 1995). Interestingly, Kubo et al. (1997) reported organ-specific susceptibility to lipid peroxidation and organ-specific differences for antioxidants, and suggested that the kidney, in terms of its fatty acid composition, is more resistant to dietary n-3 fatty acid influences. Fernandes et al. (1996) have also shown that feeding fish oil to the NZB \times NZW F₁ mouse increases programmed cell death of lymphocytes, which may prevent the accumulation of self-reactive immune cells in lymphoid organs.

Evidence for beneficial effects of n-3 fish oil fatty acids in experimentally-induced T-cell-mediated models of autoimmune arthritis appear conflicting. In the mouse, Leslie *et al.* (1985) found fish oil to protect against experimental collagen-induced autoimmune arthritis, whilst in the rat fish oil was reported to augment the disease (Prickett *et al.* 1984). We found that feeding fish oil augmented EAE in the rat (Harbige, 1993), although suppression of EAE using lower doses of fish oil has been described by Mertin (1983), but the animals developed more severe clinical disease after discontinuation of treatment. The mechanism(s) involved in fish oil-induced augmentation of experimental autoimmune disease in rats is not well understood.

Autoimmunity, immune functions and *n*-6 fatty acids

Animal studies

Erickson et al. (1983) concluded from data obtained using a mixed lymphocyte reaction (alloantigens) system as a measure of lymphocyte function in mice that high dietary levels of safflower oil (rich in linoleic acid) suppressed the response whereas lower levels intensified the response. Similarly Yaqoob et al. (1994) reported that high levels of dietary linoleic acid-rich oil suppressed Con A-stimulated lymphocyte proliferation in rats. In contrast, De Deckere et al. (1988) found no effect on spleen lymphocyte proliferation in response to Con A in rabbits and rats fed on diets containing a high amount of linoleic acid (35 % digestible energy as sunflower oil) compared with palm oil. Studies by Kelley et al. (1988) in rabbits suggest that a low intake of linoleic acid-rich safflower oil does not affect spleen-derived lymphocyte responses or peripheral-blood-derived lymphocyte responses to PHA and Con A. Cantillon et al. (1990) also reported no effect of feeding linoleic acid-rich oil on rat lymphocyte proliferation ex vivo. Spleen lymphocytes from rats orally administered safflower-seed oil showed enhanced Con A responses ex vivo compared with lymphocytes from rats fed on the basal diet (Harbige et al. 1995). Kollmorgen et al. (1979) reported that feeding rats on maize oil reduced the ex vivo lymphocyte proliferative response to Con A; an important consideration in the Kollmorgen et al. (1979) study is the

use of maize oil stripped of vitamin E. Young et al. (1987) showed reduced spleen lymphocyte proliferation in response to Con A in mice treated with a pure linoleic acid preparation at an estimated ten times higher dose than those used by other workers; in addition, linoleic acid was administered by subcutaneous injection. Cinader et al. (1983) found loss of 'suppressor activity' to the tolerance inducibility of rabbit y-globulin in mice fed on a linoleic acid-rich diet. In mice fed a y-linolenic acid-rich diet high- and low-dose oral ova tolerance induction reduced ova-specific proliferative responses (but not in animals fed on borage oil only) in ova immunized mice and decreased (high dose oral ova only) ova-specific serum IgG_{2a} and IgG_{2b} (LS Harbige and BAC Fisher, unpublished results). In summary, the effects of dietary linoleic acid on immune functions in animals appears inconsistent, and more information is required on other fatty acids of the *n*-6 family.

Human studies

The effects of n-6 fatty acids on immune functions, particularly the desaturated and elongated metabolites of linoleic acid, have not been studied as much as the n-3 fatty acids in human subjects. Supplementation with γ -linolenic acid-rich borage oil (Borago officinalis) did not affect the lymphoproliferative response to PHA and anti-CD3, but did increase the production of mitogen-stimulated peripheral-blood mononuclear cell TGF β_1 and decreased the production of IL-4 and IL-10 (Harbige & Fisher, 1997). These findings are consistent with the effects of prostaglandin (PG) E_1 and PGE₂ on IL-4 and IL-10 in human T-cell clones (Harbige et al. 1997a). However, the effects of PGE on T-cell clones reported by Harbige et al. (1997a) are not consistent with previous findings obtained using murine and human T-cell clones (Betz & Fox, 1991; Watanabe et al. 1994), possibly due to differences in clones, mode of activation and eicosanoid concentrations used. Membrane lipid-oxidation products have been shown to increase $TGF\beta_1$ expression and production (Leonarduzzi et al. 1997). It is possible, therefore, that an increase in membrane unsaturation, leading to an increased peroxidizability index, by feeding γ -linolenic acid-rich borage oil might lead to increased production of TGF β_1 . However, we found no change in plasma malondialdehyde concentration, a marker of peroxidation, in a γ -linolenic acid-rich borage oil supplementation study in human subjects (LS Harbige, F Kelley, C Dunster and K Ghebremeskel, unpublished results).

Autoimmune disease

In the NZB × NZW F₁ autoimmune mouse, linoleic acid-rich oil appears to exacerbate the disease (Levy *et al.* 1982) or to have no effect (Hurd et al. 1981). Similarly, in the MLR/Ipr autoimmune-prone mouse, linoleic acid-rich oil also appears to enhance the disease or to have no effect (Kelley *et al.* 1985; Godfrey *et al.* 1986). In addition, in the MLR/Ipr autoimmune-prone mouse, Godfrey *et al.* (1986) observed a greater survival rate for mice supplemented with 50 g evening primrose (*Oenothera biennis*) oil/kg diet (72 g linoleic acid and 9 g γ -linolenic acid, the parent *n*-6 fatty

acid, is unable to ameliorate antibody-mediated spontaneous autoimmune disease, whereas the desaturated n-6 metabolites can. Studies in the guinea-pig have shown linoleic acid to partially suppress the incidence and severity of EAE (Meade et al. 1978). Similarly, high levels of a linoleic-rich oil containing low levels of y-linolenic acid (linoleic acid : γ -linolenic acid 7 : 1, based on total fatty acid content) partially suppressed the incidence and severity of EAE in the rat (Mertin & Stackpoole, 1978). Using γ-linolenic acid-rich oils from fungal or plant sources we demonstrated complete protection against EAE in both rats and mice (Harbige et al. 1995, 1997b). This series of investigations demonstrated important disease-modifying effects of linoleic and y-linolenic acid on clinical and histopathological manifestations of EAE. Depending on dose, y-linolenic acid is completely protective in acute rat EAE, whereas linoleic acid has a dose-dependent action on the clinical severity of EAE, although not abolishing it. Analysis of spleen-cellmembrane fatty acid composition suggested that the previously described clinical response in EAE is due to conversion of γ -linolenic acid to the longer-chain *n*-6 eicosanoid precursor fatty acids, dihomo-y-linolenic and arachidonic acids.

In chronic relapsing EAE induced in SJL mice with an encephalitogenic peptide (92-106) of myelin oligodendrocyte glycoprotein, feeding γ -linolenic acid markedly inhibited both the acute and relapse phases of the disease (Harbige et al. 1997b). Spleen-cell lymphoproliferative responses to Con A, myelin oligodendrocyte glycoprotein peptide (92-106) and protein purified derivative were unaltered when comparing γ -linolenic acid-fed and control EAE mice, indicating no gross immunosuppression by the fatty acid treatment (Harbige et al. 1997b). However, on activation with myelin oligodendrocyte glycoprotein peptide, Con A, and protein purified derivative there was a nonspecific increase in the production of $TGF\beta_1$ and PGE_2 in γ -linolenic acid-fed mice ex vivo, with no effect on the production of IL-2 and interferon- γ (Harbige *et al.* 1997*b*). Furthermore, there was higher $TGF\beta_1$ mRNA expression and increased membrane eicosanoid precursor fatty acids (PGE1 and PGE₂) in the spleens of γ -linolenic acid-fed mice. These findings are consistent with reports that administration of TGFB protects in acute and relapsing EAE (Racke et al. 1993; Santambrogio et al. 1993), and that PG inhibitors, such as indomethacin, augment EAE (Ovadia & Paterson, 1982). In addition, during the natural recovery phase from EAE, TGFβ-secreting T-cells can inhibit EAE effector cells, TGFβ is expressed in the central nervous system and, in oral-tolerance-induced protection in EAE, TGF β and PGE₂ are expressed in the brain (Karpus & Swanborg, 1991; Khoury et al. 1992). Feeding γ -linolenic acid can increase PGE₁ and PGE₂ production (Fan & Chapkin, 1992; Harbige et al. 1997b), and PGE₂ has been shown to inhibit the production of Th₁ but not Th₂ cytokines in vitro (Phipps et al. 1991). Importantly, adoptive transfer experiments have shown EAE to be a T-cell-mediated autoimmune disease in which Th₁-type cytokine-producing cells cause pathology (Liblau et al. 1995). On this basis we could propose that dietary y-linolenic acid prevents T-cell-mediated autoimmune disease by altering the balance of cytokines from a Th_{1} - to a Th₂-type response. However, our findings suggest that the effects of dietary γ -linolenic acid on EAE are mediated through Th₃-like mechanisms involving TGF β_1 (Harbige *et al.* 1997*b*). It is interesting to note that Vidard *et al.* (1995) suggested from studies of T-cell tolerance and site-specific lymphokine profiles that the type of Th subset cytokine pattern produced depends on the environment in which they are found. In relation to the observations made by Pond (1996) mentioned earlier, this could be one mechanism by which local fatty acids affect Th subset cytokines.

The antioxidant enzymes superoxide dismutase (*EC* 1.15.1.1) and catalase (*EC* 1.11.1.6) have been shown to beneficially modify the clinical course of EAE (Guy *et al.* 1989). Previously we have shown that feeding a γ -linolenic acid-rich oil to rats increases the activity of superoxide dismutase in rat tissues (Phylactos *et al.* 1994). Increased antioxidant activity may therefore be an important mechanism by which linoleic acid metabolites control EAE. Similarly, Fernandes (1994) has suggested this as a mechanism in fish oil protective effects in NZB × NZW F₁ mice.

The nature of the autoimmune model appears to be important. The Th₂ cytokines IL-4, IL-6 and IL-10 are pro-inflammmatory in the NZB × NZW F₁ model, which is antibody-mediated (Fernandes, 1994). In contrast, in EAE, which is T-cell-mediated, IL-4 and IL-10 are associated with remission (Kennedy *et al.* 1992; Khoury *et al.* 1992; Liblau *et al.* 1995). Consistent with the previous findings, in the antibody-mediated NZB × NZW F₁ autoimmune model, animals fed on a high-fat diet consisting of equal amounts of lard and soyabean oil (linoleic acid-rich) develop a more severe disease and have a shortened lifespan, which is associated with increased autoantibody, and a decreased Th₁ and increased Th₂ cytokine profile (Lin *et al.* 1996).

Supplementation with *n*-6 and *n*-3 fatty acids in human autoimmune inflammatory disease

In clinical studies significant benefits have been reported in patients with systemic lupus erythematosus following a low-fat diet plus an n-3 fatty acid-rich fish oil supplement (Walton et al. 1991). Controlled trials in patients with rheumatoid arthritis on diets high in n-3 fatty acids without background saturated fat manipulation have also shown clinically beneficial effects (Kremer et al. 1985; Cleland et al. 1988). Interestingly, Kremer et al. (1987) found a significant correlation between decreases in neutrophil LTB4 production and decreases in the number of tender joints in individual patients supplemented with fish oil. In patients with rheumatoid arthritis, treatment with evening primrose oil enabled 73 % of patients to stop or reduce non-steroidal antiinflammatory drug treatment (Belch et al. 1988). Using γ -linolenic acid-rich borage oil (23 g γ -linolenic acid and 62 g linolenic acid/100 g total fatty acids), Leventhal et al. (1993) managed to significantly reduce clinically-important signs and symptoms of disease activity in patients with active rheumatoid arthritis. Belluzzi et al. (1996), feeding a novel fish oil preparation in a double-blind placebo-controlled trial with patients with Crohn's disease have reported a 60 %reduction in relapse rate. Double-blind trials to determine the therapeutic efficacy of diets supplemented with sunflower

oil, a source of linoleic acid, have been carried out in patients with multiple sclerosis by Miller et al. (1973), Bates et al. (1978) and Paty et al. (1978). In the first two studies, relapse rate and severity of disease were reduced in the treated groups, but Paty et al. (1978) showed no such effect. In a meta-analysis of the previously described trials, Dworkin et al. (1984) showed reduced relapse rate and severity, and a decrease in the long-term progression of the disease in patients with mild multiple sclerosis. Furthermore, open studies with patients with multiple sclerosis suggest that a low-fat diet and/or manipulation of dietary n-6 and n-3 fatty acids may be beneficial (Swank & Grimsgaard, 1988; Harbige et al. 1990). Importantly, many trial designs, including those for multiple sclerosis, have utilized olive oil as the placebo control and have not taken into account the total fat and saturated fat intake. It is apparent that olive oil (65–85 g oleic acid/100 g total fatty acids) increases the survival rate of MLR/Ipr mice and reduces the incidence of EAE in the guinea-pig (Meade et al. 1978; Godfrey et al. 1986). Patients with rheumatoid arthritis in the olive oil comparative control group had improvement in some clinical indices in the fish oil study of Cleland et al. (1988). Olive oil has been observed also to be of some clinical benefit in rheumatoid arthritis, with corresponding improvements in several laboratory variables, including C-reactive protein (G Darlington, personal communication). There is a clear need, therefore, for more carefully designed and controlled trials in the therapeutic application of fatty acids to human autoimmune inflammatory conditions. Since side effects are a problem with current treatment of human autoimmune disease, safe and more effective treatments would greatly improve their management. Nutritional approaches using fatty acid supplementation in human autoimmune disease, whether as an alternative or adjunct therapy, is therefore potentially very important.

Summary and conclusions

Clearly there is much evidence to show that under wellcontrolled laboratory and dietary conditions fatty acid intake can have profound effects on animal models of autoimmune disease. Studies in human autoimmune disease have been less dramatic; however, human trials have been subject to uncontrolled dietary and genetic backgrounds, infection and other environmental influences, and basic trial designs have been inadequate. The impact of dietary fatty acids on animal autoimmune disease models appears to depend on the animal model and the type and amount of fatty acids fed. Diets low in fat, essential fatty acid-deficient, or high in n-3 fatty acids from fish oils increase the survival and reduce disease severity in spontaneous autoantibody-mediated disease, whilst linoleic acid-rich diets appear to increase disease severity. In experimentally-induced T-cell-mediated autoimmune disease, essential fatty acid-deficient diets or diets supplemented with n-3 fatty acids appear to augment disease, whereas n-6 fatty acids prevent or reduce the severity. In contrast, in both T-cell and antibody-mediated auto- immune disease the desaturated and elongated metabolites of linoleic acid are protective. Suppression of autoantibody and T lymphocyte proliferation, apoptosis of autoreactive lymphocytes, and reduced pro-inflammatory cytokine production by

high-dose fish oils are all likely mechanisms by which n-3fatty acids ameliorate autoimmune disease. However, these could be undesirable long-term effects of high-dose fish oil which may compromise host immunity. The protective mechanism(s) of n-6 fatty acids in T-cell- mediated autoimmune disease are less clear, but may include dihomo-y-linolenic acid- and arachidonic acid-sensitive immunoregulatory circuits such as Th₁ responses, TGFB₁mediated effects and Th₃-like responses. It is often claimed that n-6 fatty acids promote autoimmune and inflammatory disease based on results obtained with linoleic acid only. It should be appreciated that linoleic acid does not reflect the functions of dihomo-y-linolenic and arachidonic acid, and that the endogenous rate of conversion of linoleic to arachidonic acid is slow (Hassam et al. 1975, 1977; Phylactos et al. 1994; Harbige et al. 1995). In addition to effects of dietary fatty acids on immunoregulation, inflammation as a consequence of immune activation in autoimmune disease may also be an important mechanism of action whereby dietary fatty acids modulate disease activity.

In conclusion, regulation of gene expression, signal transduction pathways, production of eicosanoids and cytokines, and the action of antioxidant enzymes are all mechanisms by which dietary n-6 and n-3 fatty acids may exert effects on the immune system and autoimmune disease. Probably the most significant of these mechanisms in relation to our current understanding of immunoregulation and inflammation would appear to be via fatty acid effects on cytokines. The amount, type and balance of dietary fatty acids and associated antioxidant nutrients appear to impact on the immune system to produce immune-deviation or immunosuppressive effects, and to reduce immune-mediated inflammation which will in turn affect the susceptibility to, or severity of, autoimmune disease.

Acknowledgements

I would like to thank my colleagues and collaborators, particularly Dr Sandra Amor, Dr Lorna Layward, Miss Margert Morris and Mr Benjamin Fisher. I am also grateful to Dr Graham Wallace and Dr Keb Ghebremeskel for reading the manuscript and helpful discussion, and the Henry Smith Charity and Hoffman–La Roche for financial support.

References

- Alexander NJ, Smythe NL & Jokinen MP (1987) The type of dietary fat affects the severity of autoimmune disease in NZB/NZW mice. American Journal of Pathology 127, 106–121.
- Bates D, Fawcett PRW, Shaw DA & Weightman D (1978) Polyunsaturated fatty acids in the treatment of acute remitting multiple sclerosis. *British Medical Journal* 2, 1390–1391.
- Belch JJ, Ansell D, Madhok R, O'Dowd A & Sturrock RD (1988) Effects of altering dietary essential fatty acids on requirements for non-steroidal anti-inflammatory drugs in patients with rheumatoid arthritis: a double blind placebo controlled study. Annals of the Rheumatic Diseases 47, 96–104.
- Belluzzi A, Brignola C, Campieri M, Pera A, Boschi S & Miglioli M (1996) Effect of an enteric-coated fish-oil preparation on relapses in Crohn's disease. *New England Journal of Medicine* 334, 1557–1560.

- Betz M & Fox BS (1991) Prostaglandin E_2 inhibits production of Th1 lymphokines but not of Th2 lymphokines. *Journal of Immunology* **146**, 108–113.
- Calder PC (1996) Immunomodulatory and anti-inflammatory effects of *n*-3 polyunsaturated fatty acids. *Proceedings of the Nutrition Society* **55**, 737–774.
- Cantillon DM, Morrissey PA, O'Connor TP & Flynn A (1990) Effect of dietary fish oil on rat lymphocyte proliferation in vitro. *Proceedings of the Nutrition Society* **49**, 113A.
- Chandrasekar B, Troyer DA, Venkatraman JT & Fernandes G (1995) Dietary omega-3 lipids delay the onset and progression of autoimmune lupus nephritis by inhibiting transforming growth factor beta mRNA and protein expression. *Journal of Autoimmunity* 8, 381–393.
- Cinader B, Clandinin MT, Hosokawa T & Robblee NM (1983) Dietary fat alters the fatty acid composition of lymphocyte membranes and the rate at which suppressor capacity is lost. *Immunology Letters* 6, 331–337.
- Clausen J & Moller J (1967) Allergic encephalomyelitis induced by brain antigen after deficiency in polyunsaturated fatty acids during myelination. Acta Neurologica Scandinavica 43, 375–388.
- Cleland LG, French JK, Betts WH, Murphy GA & Elliott MJ (1988) Clinical and biochemical effects of dietary fish oil supplements in rheumatoid arthritis. *Journal of Rheumatology* 15, 1471–1475.
- De Deckere EM, Verplanke CJ, Blonk CG & Van Nielen WG (1988) Effect of type and amount of dietary fat on rabbit and rat lymphocyte proliferation in vitro. *Journal of Nutrition* **118**, 11–18.
- Dvorak AM, Dvorak HF, Peters SP, Shulman ES, MacGlashan DW, Pyne K, Harvey VS, Galli SJ & Lichtenstein LM (1983) Lipid bodies: cytoplasmic organelles important to arachidonate metabolism in macrophages and mast cells. *Journal of Immunol*ogy 131, 2965–2976.
- Dworkin RH, Bates D, Millar JHD & Paty DW (1984) Linoleic acid and multiple sclerosis: a reanalysis of three double-blind trials. *Neurology* 34, 1441–1445.
- Erickson KL, Adams DA & McNeil CJ (1983) Dietary lipid modulation of immune responsiveness. *Lipids* 18, 468–474.
- Fan Y & Chapkin RS (1992) Mouse peritoneal macrophage prostaglandin E₁ synthesis is altered by dietary gamma-linolenic acid. *Journal of Nutrition* **122**, 1600–1606.
- Fernandes G (1994) Dietary lipids and risk of autoimmune disease. Clinical Immunology and Immunopathology **72**, 193–197.
- Fernandes G, Bysani C, Venkatraman JT, Tomar V & Zhao W (1994) Increased TGF- β and decreased oncogene expression by ω -3 fatty acids in the spleen delays onset of autoimmune disease in B/W mice. *Journal of Immunology* **152**, 5979–5987.
- Fernandes G, Chandrasekar B, Luan X & Troyer DA (1996) Modulation of antioxidant enzymes and programmed cell death by *n*-3 fatty acids. *Lipids* **31**, Suppl., S91–S96.
- Fernandes G, Yunis EJ, Smith J & Good RA (1972) Dietary influence on breeding behaviour, hemolytic anemia and longevity in NZB mice. Proceedings of the Society for Experimental Biology and Medicine 139, 1189–1196.
- Fujikawa M, Yamashita N, Yamazaki K, Sugiyama E, Suzuki H & Hamazaki T (1992) Eicosapentaenoic acid inhibits antigenpresenting cell function of murine splenocytes. *Immunology* 75, 330–335.
- Godfrey DG, Stimson WH, Watson RR, Belch JJ & Sturrock D (1986) The modulation of autoimmunity in the MRL-mp-Ipr/Ipr mouse by dietary fatty acid supplementation. *Progress in Lipid Research* 25, 298–293.
- Goetzl EJ, An S & Smith WL (1995) Specificity of expression and effects of eicosanoid mediators in normal physiology and human disease. *FASEB Journal* 9, 1051–1058.

- Guy J, Ellis EA, Hope GM & Rao A (1989) Antioxidant enzyme suppression of demyelination in experimental optic neuritis. *Current Eye Research* 8, 467–477.
- Harbige LS (1993) Essential fatty acids, experimental autoimmune encephalomyelitis and multiple sclerosis. PhD Thesis, Brunel University.
- Harbige LS (1996) Nutrition and immunity with emphasis on infection and autoimmune disease. *Nutrition and Health* **10**, 285–312.
- Harbige LS & Fisher BAC (1997) Effect of omega-6 lipid-rich borage oil feeding on immune function in healthy volunteers. *Transactions of the Biochemical Society* 25, 343S.
- Harbige LS, Jones R, Jenkins R, Fitzgerald G, Forti A & Budowski P (1990) Nutritional management in multiple sclerosis with reference to experimental models. Upsala Journal of Medical Science 48, 189–207.
- Harbige LS, Layward L, Morris M & Amor S (1997a) Cytokine secretion by human T cell clones is differentially regulated by eicosanoids. *Transactions of the Biochemical Society* 25, 347S.
- Harbige LS, Layward L, Morris M & Amor S (1997b) Protective mechanisms by omega-6 lipids in experimental autoimmune encephalomyelitis (EAE) are associated with cytokines and eicosanoids. *Transactions of the Biochemical Society* 25, 342S.
- Harbige LS, Yeatman N, Amor S & Crawford MA (1995) Prevention of experimental autoimmune encephalomyelitis in Lewis rats by a novel fungal source of γ -linolenic acid. *British Journal of Nutrition* **74**, 701–715.
- Hassam AG, Rivers PW & Crawford MA (1977) Metabolism of gamma-linolenic acid in essential fatty acid-deficient rats. *Journal of Nutrition* **107**, 519–524.
- Hassam AG, Sinclair AJ & Crawford MA (1975) The incorporation of orally fed radioactive gamma-linolenic acid and linoleic acid into the liver and brain lipids of suckling rats. *Lipids* **10**, 417–420.
- Hennig B, Toborek M, McClain CJ & Diana JC (1996) Nutritional implications in vascular endothelial cell metabolism. *Journal of* the American College of Nutrition 15, 345–358.
- Hopkins J, McConnell I & Pearson JD (1981) Lymphocyte traffic through antigen-stimulated lymph nodes II: role of prostaglandin E_2 as a mediator of cell shutdown. *Immunology* **42**, 225–231.
- Horiguchi J, Spriggs D, Imamura K, Stone R, Luebbers R & Kufe D (1989) Role of arachidonic acid metabolism in transcriptional induction of tumor necrosis factor gene expression by phorbol ester. *Molecular and Cellular Biology* 9, 252–258.
- Huang SC, Misfeldt ML & Fritsche KL (1992) Dietary fat influences Ia antigen expression and immune cell populations in the murine peritoneum and spleen. *Journal of Nutrition* 122, 1219–1231.
- Hughes DA, Pinder AC, Piper Z & Lund EK (1995) N-3 polyunsaturated fatty acids (PUFA) modulate the expression of functionally associated molecules on human monocytes. *Transactions of the Biochemical Society* **23**, 303S.
- Hurd ER, Johnston JM, Okita JR, MacDonald PC, Ziff M & Gilliam JM (1981) Prevention of glomerulonephritis and prolonged survival in New Zealand black/New Zealand white F1 hybrid mice fed an essential fatty acid-deficient diet. *Journal of Clinical Investigation* 67, 478–485.
- Karpus WJ & Swanborg RH (1991) CD4⁺ suppressor cells inhibit the function of effector cells of experimental autoimmune encephalomyelitis through a mechanism involving transforming growth factor- β . Journal of Immunology **146**, 1163–1168.
- Kelley DS, Nelson GJ, Serrato CM, Schmidt PC & Branch LB (1988) Effects of type of dietary fat on indices of immune status in rabbits. *Journal of Nutrition* **118**, 1376–1384.
- Kelley VE, Ferretti A, Izui S & Strom TB (1985) A fish oil diet rich in eicosapentaenoic acid reduces cyclooxygenase metabolites, and suppresses lupus in MRL-Ipr mice. *Journal of Immunology* 134, 1914–1919.

- Kennedy MK, Torrance DS, Picha KS & Mohler KM (1992) Analysis of cytokine mRNA expression in the central nervous system of mice with experimental autoimmune encephalomyelitis reveals that IL-10 mRNA expression correlates with recovery. *Journal of Immunology* 149, 2496–2505.
- Khoury SJ, Hancock WW & Weiner HL (1992) Oral tolerance to myelin basic protein and natural recovery from experimental autoimmune encephalomyelitis are associated with downregulation of inflammatory cytokines and differential upregulation of transforming growth factor β , interleukin 4, and prostaglandin E expression in the brain. *Journal of Experimental Medicine* **176**, 1355–1364.
- Kremer JM, Bigaouette J, Michalek AV, Timchalk MA, Lininger L, Rynes RI, Huyck C, Zieminski J & Bartholomew LE (1985) Effects of manipulation of dietary fatty acids on clinical manifestations of rheumatoid arthritis. *Lancet* i, 184–187.
- Kremer JM, Jubiz W, Michalek A, Rynes RI, Bartholomew LE, Bigaouette J, Timchalk M, Beeler D & Lininger L (1987) Fish-oil fatty acid supplementation in active rheumatoid arthritis: a double-blinded, controlled, crossover study. *Annals of Internal Medicine* 106, 497–502.
- Kollmorgen GM, Sansing WA, Lehman AA, Fischer G, Longley RE, Alexander SS, King MM & McCay PB (1979) Inhibition of lymphocyte function in rats fed high-fat diets. *Cancer Research* 39, 3458–3462.
- Kubo K, Saito M, Tadokoro T & Maekawa A (1997) Changes in susceptibility of tissues to lipid peroxidation after ingestion of various levels of docosahexaenoic acid and vitamin E. British Journal of Nutrition 78, 655–669.
- Lefkowith J, Schreiner G, Cormier J, Handler ES, Driscoll HK, Greiner D, Mordes JP & Rossini AA (1990) Prevention of diabetes in the BB rat by essential fatty acid deficiency: relationship between physiological and biochemical changes. *Journal of Experimental Medicine* **171**, 729–742.
- Lefkowith JB (1988) Essential fatty acid deficiency inhibits the in vivo generation of leukotriene B₄ and suppresses levels of resident and elicited leukocytes in acute inflammation. *Journal of Immunology* 140, 228–233.
- Leitch GA, Lee TH, Ringel EW, Prickett JD, Robinson DR, Pyne SG, Corey EJ, Drazen JM, Austen KF & Lewis RA (1984) Immunologically induced generation of tetraene and pentaene leukotrienes in the peritoneal cavities of menhaden-fed rats. *Journal of Immunology* **132**, 2556–2565.
- Leonarduzzi G, Scavazza A, Biasi F, Chiarpotto E, Camandola S, Vogl S, Dargel R & Poli G (1997) The lipid peroxidation end product 4-hydroxy-2,3-nonenal up-regulates transforming growth factor β_1 expression in the macrophage lineage: a link between oxidative injury and fibrosclerosis. *FASEBS Journal* 11, 851–857.
- Leslie CA, Gonnerman WA, Ullman MD, Hayes KC, Franzblau C & Cathcart ES (1985) Dietary fish oil modulates macrophage fatty acids and decreases arthritis susceptibility in mice. *Journal* of Experimental Medicine 162, 1336–1349.
- Leventhal LJ, Boyce EG & Zurier RB (1993) Treatment of rheumatoid arthritis with gamma linolenic acid. Annals of Internal Medicine 119, 867–873.
- Levine S & Sowinski R (1980) Effect of essential fatty acid deficiency on experimental allergic encephalomyelitis in rats. *Journal of Nutrition* **110**, 891–896.
- Levy JA, Ibrahim AB, Shirai T, Ohta K, Nagasawa R, Yoshida H, Estes J & Gardner M (1982) Dietary fat affects immune response, production of antiviral factors, and immune complex disease in NZB/NZW mice. *Proceedings of the National Academy of Sciences USA* 79, 1974–1978.

- Liblau RS, Singer SM & McDevitt HO (1995) Th1 and Th2 CD4⁺ T cells in the pathogenesis of organ-specific autoimmune diseases. *Immunology Today* **16**, 34–38.
- Lin B, Huang C, Chiang B & Jeng S (1996) Dietary fat influences Ia antigen expression, cytokines and prostaglandin E_2 production of immune cells in autoimmune-prone NZB × NZW F1 mice. *British Journal of Nutrition* **75**, 711–722.
- Lin B, Jeng S, Chiang B & Huang C (1997) Dietary fat affects lipids and anti-cardiolipin antibody levels in autoimmune-prone NZB/W F1 mice. *British Journal of Nutrition* 77, 657–669.
- Male D, Cooke A, Owen M, Trowsdale J & Champion B (1996) Advanced Immunology, 3rd ed. London: Mosby.
- Meade CJ, Mertin J, Sheena J & Hunt R (1978) Reduction by linoleic acid of the severity of experimental allergic encephalomyelitis in the guinea-pig. *Journal of the Neurological Sciences* 35, 291–308.
- Medawar PB, Hunt R & Mertin J (1979) An influence of diet on transplantation immunity. *Proceedings of the Royal Society* B 206, 265–280.
- Mertin J (1981) Essential fatty acids and cell-mediated immunity. Progress in Lipid Research 20, 851–856.
- Mertin J (1983) Omega-6 and omega-3 polyunsaturates and the immune system. British Journal of Clinical Practice 31, Suppl., 111-114.
- Mertin J & Stackpoole A (1978) Suppression by essential fatty acids of experimental allergic encephalomyelitis is abolished by indomethacin. *Prostaglandins and Medicine* **1**, 283–291.
- Meydani SN, Lichtenstein AH, Cornwall S, Meydani M, Goldin BR, Rasmussen H, Dinarello CA & Schaefer EJ (1993) Immunologic effects of national cholesterol education panel step-2 diets with and without fish-derived *n*-3 fatty acid enrichment. *Journal* of Clinical Investigation **92**, 105–113.
- Millar JHD, Zilkha KJ, Langman MJS, Payling-Wright H, Smith AD, Belin J & Thompson RHS (1973) Double-blind trial of linoleate supplementation of the diet in multiple sclerosis. *British Medical Journal* 1, 765–768.
- Morrissey PA, O'Connor TP & Flynn A (1990) Effect of dietary fish oil on rat lymphocyte proliferation in vitro. *Proceedings of the Nutrition Society* **49**, 113A.
- Morrow JW, Ohashi Y, Hall J, Pribnow J, Hirose S, Shirai T & Levy JA (1985) Dietary fat and immune function I. Antibody responses, lymphocyte and accessory cell function in NZB \times NZW F₁ mice. *Journal of Immunology* **135**, 3857–3863.
- Mosman TR & Coffman RL (1989) Th1 and Th2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annual Review of Immunology* 7, 145–173.
- Ovadia H & Paterson PY (1982) Effect of indomethacin treatment upon actively-induced and transferred experimental allergic encephalomyelitis (EAE) in Lewis rats. *Clinical and Experimental Immunology* **49**, 386–392.
- Paty DW, Cousin HK, Read S & Adlakha K (1978) Linoleic acid in multiple sclerosis: failure to show any therapeutic benefit. *Acta Neurologica Scandinavica* **58**, 53–58.
- Phipps RP, Stein SH & Roper RL (1991) A new view of prostaglandin E regulation of the immune response. *Immunology Today* 12, 349–352.
- Phylactos AC, Harbige LS & Crawford MA (1994) Essential fatty acids alter the activity of manganese-superoxide dismutase in rat heart. *Lipids* 29, 111–115.
- Pond CM (1996) Interactions between adipose tissue and the immune system. *Proceedings of the Nutrition Society* 55, 111–126.
- Prickett JD, Trentham DE & Robinson DR (1984) Dietary fish oil augments the induction of arthritis in rats immunized with type II collagen. *Journal of Immunology* **132**, 725–729.

- Pushkareva M, Obeid LM & Hannun YA (1995) Ceramide: an endogenous regulator of apoptosis and growth suppression. *Immunology Today* 16, 294–302.
- Racke MK, Sriram S, Carllini J, Cannella B, Raine CS & McFarlin DE (1993) Long-term treatment of chronic relapsing experimental allergic encephalomyelitis by transforming growth factor- $\beta 2$. Journal of Neuroimmunology **46**, 175–183.
- Robinson DR, Colvin RB, Hirai A & Steinberg AD (1986) Dietary marine lipids modify autoimmune diseases. In *Health Effects* of Polyunsaturated Fatty Acids in Seafoods, pp. 247–259 [A Simopoulos, RR Kifer and RE Martin, editors]. New York: Academic Press Inc.
- Ross R (1993) The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature* 362, 801–809.
- Roper LR, Conrad DH, Brown DM, Warner GL & Phipps R (1990) Prostaglandin E2 promotes IL-4-induced IgE and IgG1 synthesis. *Journal of Immunology* 145, 2644–2651.
- Santambrogio L, Hochwald GM, Saxena B, Leu CH, Martz JE, Carlino JA, Ruddle NH, Palladino MA, Gold LI & Thorbecke GJ (1993) Studies on the mechanisms by which transforming growth factor-β protects against allergic encephalomyelitis. *Journal of Immunology* **151**, 1116–1127.
- Schlager SI, Meltzer MS & Madden LD (1983) Role of membrane lipids in the immunological killing of tumor cells: II. Effector cell lipids. *Lipids* 18, 483–488.
- Schmidt EB & Dyerberg J (1989) n-3 Fatty acids and leucocytes. Journal of Internal Medicine 225, Suppl. 1, 151–158.
- Selivonchick DP & Johnston PV (1975) Fat deficiency in rats during development of the central nervous system and susceptibility to experimental allergic encephalomyelitis. *Journal of Nutrition* 105, 288–300.
- Sherrington EJ, Sanderson P & Calder PC (1995) The effect of dietary lipid manipulation on macrophage cell surface expression. *Transactions of the Biochemical Society* 23, 272S.
- Shipman PM, Schmidt RR & Chepenik KP (1988) Relation between arachidonic acid metabolism and development of thymocytes in fetal thymic organ cultures. *Journal of Immunol*ogy 140, 2714–2720.
- Swank RL & Grimsgaard A (1988) Multiple sclerosis: the lipid relationship. American Journal of Clinical Nutrition 48, 1387–1393.

- Tate G, Mandell BF, Laposata M, Ohliger D, Baker DG, Schumacher RH & Zurier RB (1989) Suppression of acute and chronic inflammation by dietary gamma-linolenic acid. *Journal* of *Rheumatology* **16**, 729–733.
- Terano T, Salmon JA & Moncada S (1984) Effect of orally administered eicosapentaenoic acid (EPA) on the formation of leukotriene B_4 and leukotriene B_5 by rat leukocytes. *Biochemical Pharmacology* **33**, 3071–3076.
- Vidard L, Colarusso LJ & Benacerraf B (1995) Specific T-cell tolerance may reflect selective activation of lymphokine synthesis. *Proceedings of the National Academy of Sciences USA* 92, 2259–2262.
- Walton AJE, Snaith ML, Locnistar M, Cumberland AG, Morrow WJW & Isenberg DA (1991) Dietary fish oil and the severity of symptoms in patients with systemic lupus erythematosus. *Annals of the Rheumatic Diseases* 50, 463-466.
- Watanabe S, Yssel H, Harada Y & Arai K (1994) Effects of prostaglandin E_2 on Th0-type human T cell clones: modulation of functions of nuclear proteins involved in cytokine production. *International Immunology* **6**, 523–532.
- Weiner HL (1997) Oral tolerance: immune mechanisms and the treatment of autoimmune diseases. *Immunology Today* 18, 335–343.
- Wu D, Meydani SN, Meydani M, Hayek MG, Huth P & Nicolosi RJ (1996) Immunologic effects of marine- and plant-derived n-3 polyunsaturated fatty acids in nonhuman primates. *American Journal of Clinical Nutrition* 63, 273–280.
- Yaqoob P & Calder PC (1995) The effect of dietary lipid manipulation on the production of murine T cell-derived cytokines. *Cytokine* 7, 548–553.
- Yaqoob P, Newsholme EA & Calder PC (1994) The effect of dietary lipid manipulation on rat lymphocyte subsets and proliferation. *Immunology* **82**, 603–610.
- Young MR, Ellis NK, Young ME & Wepsic HT (1987) Stimulation of hematopoiesis and bone marrow suppressor cells by the subcutaneous injection of linoleic acid. *Cellular Immunology* **107**, 238–248.
- Yumura W, Hattori S, Morrow WJ, Mayes DC, Levy JA & Shirai T (1985) Dietary fat and immune function. *Journal of Immunology* 135, 3864–3868.

© Nutrition Society 1998